

Table 4 Mean performance for cane yield and milliblecane of 16 sugarcane genotypes evaluated at two locations of Khyber Pakhtunkhwa during 2011-12 and 2012-13

Genotypes	Cane yield (t/ha)			Millable cane		
	Mardan	Harichand	Genotype Mean	Mardan	Harichand	Genotype Mean
MS91CP272	65.10	51.62	58.36	78.17	69.67	73.92
MS94CP15	70.92	52.34	61.63	70.00	63.17	66.58
MS91CP238	67.52	52.29	59.90	91.67	80.67	86.17
MS92CP979	73.16	51.48	62.32	81.33	94.33	87.83
MS99HO391	66.99	52.91	59.95	87.67	66.50	77.08
S97CP288	71.62	33.91	52.76	85.17	60.17	72.67
MS99HO317	88.98	47.86	68.42	91.67	69.33	80.50
RS97N45	75.13	38.96	57.04	74.00	72.67	73.33
MS99HO388	70.45	35.29	52.87	87.17	54.50	70.83
MS99HO675	66.59	45.86	56.22	80.50	71.17	75.83
MS99HO93	60.47	32.00	46.24	89.83	47.83	68.83
S96SP1215	65.48	30.72	48.10	74.00	43.50	58.75
Hoth127	59.20	42.05	50.63	80.33	70.50	75.42
CP89831	65.72	50.91	58.31	83.34	81.50	82.42
CP77400	66.97	48.10	57.53	86.17	80.67	83.42
Mardan93	65.03	43.67	54.35	76.50	80.67	78.58
Location Mean	68.71	44.37		78.17	69.67	
Genotype LSD (0.05)		10.77			Ns	
Location LSD (0.05)		3.81			Ns	
G X L LSD (0.05)		15.23			Ns	

Note: Ns = Non-significant, G X L = Genotype x loca