

Table 2 Means of Pre-emergence (T0-T4) and Post-emergence (T5-T7) herbicide treatments on various weeds in Gladiolus filed

Treatments	Cyperus rotundus	Carvensis	Coronopus didymus	Chenopodium album	Anagallis arvensis	Conyza stricta	Stellaria media	Poa annua	Melilotus didymus
T ₀ (Control)	25.11	3.88	4.00	10.55	12.77	6.33	12.00	10.22	4.44
T ₁ (Metribuzine)	19.44	1.00	0.44	0.55	0.55	1.00	0.44	0.33	0.66
T ₂ (S-metolachlore)	2.22	2.00	0.55	0.33	1.22	2.00	0.33	0.22	0.44
T ₃ (Pendimethalin low)	17.77	1.33	1.11	0.33	1.00	1.66	0.33	0.00	0.55
T ₄ (Pendimethalin high)	15.22	0.88	0.33	0.22	0.22	0.88	0.22	0.00	0.22
T ₅ (Paraquat)	6.00	1.33	1.11	2.44	4.11	1.00	2.44	2.44	2.33
T ₆ (Glyphosate)	3.55	1.00	1.33	1.22	3.33	0.88	2.11	2.11	1.11
T ₇ (Metribuzine low)	23.44	3.00	0.33	0.33	0.22	1.00	0.77	0.22	0.66

Note: Where, T₀=Control, T₁=Metribuzine, T₂=S-metolachlore, T₃=Pendimethalin low, T₄=Pendimethalin high, T₅=Paraquat, T₆=Glyphosate, and T₇=Metribuzine low