

Table 3 Effect of high concentrations of *C. citratus* ethanol extracts on the mortality of 0-24 h old adult *S. zeamias*

Concentratio n (%)	% mortality after						
	1 day	2 days	3 days	4 days	5 days	6 days	7 days
Control 1	0.00 ± 0.0	0.00±0.0	0.00 ± 0.0	0.00± 0.0	0.00 ± 0.0	0.00± 0.0	0.00± 0.0
Control 2	0.00 ± 0.0	0.00±0.0	0.00 ± 0.0	0.00± 0.0	0.00 ± 0.0	0.00± 0.0	0.00± 0.0
5	0.00 ± 0.0	0.00±0.0	0.00 ± 0.0	0.00± 0.0	0.00 ± 0.0	0.00± 0.0	0.00± 0.0
10	0.00 ± 0.0	0.00±0.0	0.00 ± 0.0	0.00± 0.0	0.00 ± 0.0	0.00± 0.0	0.00± 0.0
15	0.00 ± 0.0	0.00±0.0	0.00 ± 0.0	0.00± 0.0	0.00 ± 0.0	0.00± 0.0	0.00± 0.0
20	0.00 ± 0.0	0.00±0.0	0.00 ± 0.0	0.00± 0.3	0.00 ± 0.3	0.00± 0.3	0.00± 0.3
25	0.00 ± 0.0	0.00±0.0	0.00 ± 0.3	0.00± 0.3	0.00 ± 0.3	0.00± 0.3	0.00± 0.3

Note: Each value is mean ± standard error of three replicates; Values followed by the same letter in the same column are not significantly different at P>0.05 using New Duncan's Multiple Range Test; Control 1: grains without extract and extracting solvent; Control 2: grains without extracts but with extracting solvent (ethanol)